

obstacle. This approach can be considered as an integral one for the study of a system at the moment of measurements, as a single whole intact nondestructive unity with the minimum interference. It can demonstrate an awareness of potential applications for polymer science, bio- and nano- technology, medicine and for environment protection.

References

- [1] V.J. Klenin, *Thermodynamics of Systems Containing Flexible Chain Polymers* (Elsevier, 1999).
- [2] K.S. Shifrin, G. Tonna // *in Advances in Geophysics*, Editors Renata Dmowska, Barry Saltzman, Academic Press, **34**, pp. 175 - 252 (1993).
- [3] A.G. Bezrukova // *Proceedings of SPIE* **3107** (1997) 298.
- [4] A.G. Bezrukova // *Proceedings of Material Research Society* **711** (2002) paper FF7.9, p. 1.
- [5] O.L. Vlasova and A.G. Bezrukova // *Proceedings of SPIE* **5127** (2003) 154.
- [6] A.G. Bezrukova // *Proceedings of SPIE* **5400** (2004) 189.
- [7]. A.G. Bezrukova // CD: *2006 Spring National Meeting Conference Proceedings*, New York: AIChE, 2006, 6 pages.
- [8]. A.G. Bezrukova // *Proceedings of SPIE* **6253** (2006) 62530C-1.
- [9] A. Bezrukova, M. Lubomska, P. Magri, M. Rogalski // *Proceedings of SPIE* **6597** (2007) 65970M-1.
- [10] A. Bezrukova, M. Lubomska, M. Rogalski // *Rev. Adv. Mater. Sci.* **20** (2009) 70.
- [11] M. Malmsten, B. Lassen // *in Proteins at Interfaces II*, edited by T.A. Horbert and J.L. Brash, Washington DC: American Chemical Society, 1995: 228-238.
- [12] F.M. Goltsman, *Physical Experiment and Statistical Conclusions*. Leningrad, 1982, in Russian.